

Monkeypox (Mpox)



Get the vaccine to protect yourself and others

Red Door is offering testing, treatment, and vaccine.



For more information or to
schedule an appointment

612-543-5555

reddoorclinic.org/monkeypox



RED DOOR
Sexual health care for all people

About the vaccine



- It is a two-dose series; 28 days apart.
- The vaccine can be given in the forearm, shoulder, or upper back near the shoulder blade.
- Eligibility has expanded and Red Door has same-day appointments.

Who can get the vaccine



- People exposed to Mpox
- People living with HIV
- Men who have sex with men
- Transgender people
- Others who are at high risk

After care



- Your injection site might be red, swollen, and itchy.
- Loosely cover with a bandage to keep from scratching.



Things you can do to protect yourself and partners against Mpox

Consider changing some behaviors until
two weeks after your second dose of vaccine

Talk with your partners



- Exchange contact information
- Discuss what you and your partners are comfortable with for prevention steps
- Ask how they're feeling and if they have any new symptoms

Limit skin-to-skin contact



- Wear clothes while having sex and at sex parties
- Masturbate together or have phone sex
- Use condoms, gloves, and dental dams

If you think you have Mpox

Avoid these things



- Sex of any kind (oral, anal, vaginal)
- Touching each other's bodies and rash
- Kissing or exchanging spit
- Sharing towels, bedding, toothbrushes, fetish gear, and sex toys

Red Door recommends condom use, withdrawal (pulling out), and non-penetrative sex for 8 to 12 weeks after the Mpox rash has healed completely.